



Public Health
England

NHS



COVID-19 vaccination

A guide for children and young people



Find out more at [nhs/CovidVaccine](https://nhs.uk/CovidVaccine)

This leaflet explains the coronavirus (COVID-19) vaccination programme for eligible children and young people.

What is COVID-19 or coronavirus?

COVID-19 is a very infectious respiratory disease caused by the SARS-CoV-2 virus. Very few children and young people with COVID-19 infection go on to have severe disease.

There is no cure for COVID-19 although some newly tested treatments do help to reduce the risk of complications.



The NHS is offering COVID-19 vaccine to children and young people. This includes those aged 12 to 17 years at increased risk from infection who will need 2 doses of the vaccine 8 weeks apart. All other young people aged 12 to 17 years will be offered a first dose of vaccine. The timing of a second dose for these 12 to 17 year olds will be confirmed later.

Are you at risk from COVID-19 infection?

Coronavirus can affect anyone. Some children and young people are at greater risk including those living with serious conditions such as:

- cancers (such as leukaemia or lymphoma)
- diabetes
- serious heart problems
- chest complaint or breathing difficulties, including poorly controlled asthma
- kidney, liver or a gut disease
- lowered immunity due to disease or treatment (steroid medication, chemotherapy or radiotherapy)
- an organ transplant
- a neurodisability or neuromuscular condition
- a severe or profound learning disability
- Down's syndrome
- a problem with your spleen, eg sickle cell disease, or you have had your spleen removed
- epilepsy
- serious genetic problems
- other serious medical conditions as advised by your doctor or specialist.

Your specialist or GP will tell you if you need the COVID-19 vaccination.

For most children and young people COVID-19 is usually a milder illness that rarely leads to complications. For a very few the symptoms may last for longer than the usual 2 to 3 weeks. The vaccination will help to protect you against COVID-19.

Currently the vaccine licensed for children and young people is the Pfizer vaccine. This is what you will be offered.

Will the vaccine protect me?

The COVID-19 vaccination will reduce the chance of you suffering from COVID-19 disease. It may take a few weeks for your body to build up some protection from the vaccine. You should get good protection from the first dose, having the second dose should give you longer lasting protection against the virus.

Like all medicines, no vaccine is completely effective – some people may still get COVID-19 despite having a vaccination, but this should be less severe. Further information is available on symptoms on [NHS.UK](https://www.nhs.uk).

The vaccines do not contain organisms that grow in the body, and so are safe for children and young people with disorders of the immune system. These people may not respond so well to the vaccine but it should offer them protection against severe disease.

What about giving consent?

You will probably want to share information about the vaccine with your parents and discuss it together.

If you are being offered the vaccination at school, you may be given a consent form that your parent/guardian should sign giving permission for you to have the vaccination.

The nurse or GP will discuss the COVID-19 vaccine with you at your appointment and will be able to answer any questions you may have.

Common side effects

Like all medicines, vaccines can cause side effects. Most of these are mild and short term, and not everyone gets them. With the vaccine we use in under-18s, side effects are more common with the second dose.

Very common side effects include:

- having a painful, heavy feeling and tenderness in the arm where you had your injection. This tends to be worst around 1 to 2 days after the vaccination
- feeling tired
- headache
- general aches, or mild flu like symptoms

You can rest and take paracetamol (follow the dose advice in the packaging) to help make you feel better. Although feeling feverish is not uncommon for 2 to 3 days, a high temperature is unusual and may indicate you have COVID-19 or another infection. Symptoms following vaccination normally last less than a week. If your symptoms seem to get worse or if you are concerned, you or your parents can call NHS 111.


Less common side effects

Recently, cases of inflammation of the heart (called myocarditis or pericarditis) have been reported very rarely after COVID-19 vaccines.

Most of these cases have been in younger men and usually a few days after the second vaccination. Most people recovered and felt better following rest and simple treatments.

You should seek medical advice urgently if you experience:


chest pain



shortness of breath



feelings of having a fast-beating, fluttering, or pounding heart



Keep your record card safe

If you or your parents or carers do seek advice from a doctor or nurse, make sure you tell them about your vaccination (show them the vaccination card, if possible) so that they can assess you properly.

Don't forget your
COVID-19 vaccination

NHS

Make sure you keep this record
card in your purse or wallet

For more information on the
COVID-19 vaccination or what
to do after your vaccination,
see www.nhs.uk/covidvaccine

COVID-19 immunisation
Protect yourself.



You or your parents and carers can also report suspected side effects to vaccines and medicines online through the Yellow Card scheme.



[coronavirus-yellowcard.
mhra.gov.uk](https://coronavirus-yellowcard.mhra.gov.uk)

Can you catch COVID-19 from the vaccine?

You cannot catch COVID-19 from the vaccine but it is possible to have caught COVID-19 and not have the symptoms until after your vaccination appointment. The most important symptoms of COVID-19 are recent onset of any of the following:

- a new continuous cough
- a high temperature
- a loss of, or change in, your normal sense of taste or smell

If you have the symptoms above, stay at home and arrange to have a test.

What to do next

After you have had the first dose you may be given a second appointment sometime later. Your record card will show the details of the first dose.

You will be advised on the right timing for your second dose to help give the best, and longest lasting protection for you.

If you are not well when your appointment is due

You should not attend a vaccine appointment if you are self-isolating, waiting for a COVID-19 test or within 4 weeks of having a positive COVID-19 test.

How is COVID-19 spread?

COVID-19 is spread through droplets breathed out from the nose or mouth, particularly when speaking or coughing.

It can also be picked up by touching your eyes, nose and mouth after contact with contaminated objects and surfaces.



You MUST still follow any national or local restrictions and:

where advised wear a face mask



wash your hands regularly



open windows to let fresh air in



follow the current guidance



www.gov.uk/coronavirus

Further information

You or your parents should read the product information leaflet for more details on your vaccine, including possible side effects, on the Coronavirus Yellow Card website.

You can also report suspected side effects on the same website or by downloading the Yellow Card app: coronavirus-yellowcard.mhra.gov.uk/productinformation

Further information is available from nhs.uk/conditions/coronavirus-covid-19/coronavirus-vaccination/



© Crown copyright 2021. Public Health England gateway number: 2021504.
Product code: COV2021CYPEN. 3.5M 1p AUG 2021 (APS). Version 1. 24/08/2021.

If you want to order more copies of this leaflet using the product code COV2021CYPEN, please visit: www.healthpublications.gov.uk